

Online Research @ Cardiff

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/104614/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Elmaghrabi, Adel M., Rogers, Hilary ORCID: <https://orcid.org/0000-0003-3830-5857>, Francis, Dennis and Ochatt, Sergio J. 2017. PEG induces high expression of the cell cycle checkpoint gene WEE1 in embryogenic callus of *Medicago truncatula*: potential link between cell cycle checkpoint regulation and osmotic stress. *Frontiers in Plant Science* 8 , 1479. 10.3389/fpls.2017.01479 file

Publishers page: <http://dx.doi.org/10.3389/fpls.2017.01479>
<<http://dx.doi.org/10.3389/fpls.2017.01479>>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies.

See

<http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



PEG induces high expression of the cell cycle checkpoint gene *WEE1* in embryogenic callus of *Medicago truncatula*: potential link between cell cycle checkpoint regulation and osmotic stress

Adel M. Elmaghrabi^{1,2}, Hilary J. Rogers², Dennis Francis², Sergio J. Ochart^{3*}

¹ *Biotechnology Research Center (BTRC), Box 30313, Tripoli-LIBYA.*

² *School of Biosciences, Cardiff University, Park Place, Cardiff CF10 3TL, U.K.*

^{3*} *Agroécologie, AgroSup Dijon, INRA, Univ. Bourgogne Franche-Comté, F-21000 Dijon, France.*

Corresponding Author

Author emails:

Adel Elmaghrabi maghrabiam@hotmail.com

Hilary Rogers rogershj@cf.ac.uk

Dennis Francis dfrancis5@sky.com

Sergio Ochart sergio.ochatt@inra.fr

Abstract

Polyethylene glycol (PEG) can be used to mimic osmotic stress in plant tissue cultures to study mechanisms of tolerance. The aim of this experiment was to investigate the effects of PEG (M.W. 6000) on embryogenic callus of *Medicago truncatula*. Leaf explants were cultured on MS medium with 2 mg L⁻¹ NAA and 0.5 mg L⁻¹ BAP for 5 months. Then, calli were transferred to the same medium further supplemented with 10% (w/v) 6000 PEG for six months in order to study physiological and putative molecular markers of water stress. There were no significant differences in growth rate of callus or mitotic index \pm PEG although embryogenic potential of PEG treated callus was morphologically enhanced. Cells were rounder on PEG medium and cell size, nuclear size and endoreduplication increased in response to the PEG treatment. Significant increases in soluble sugar and proline accumulation occurred under PEG treatment compared with the control. Significantly, high *MtWEE1* and *MtCCS52* expression resulted from 6 months of PEG treatment with no significant differences in *MtSERK1* or *MtP5CS* expression but down regulation of *MtSOS* expression. The results are consistent in showing elevated expression of a cell cycle checkpoint gene, WEE1. It is likely that the cell cycle checkpoint surveillance machinery, that would include *WEE1* expression, is ameliorating the effects of the stress imposed by PEG.

217 words

Keywords: cell cycle, cell division, cell morphology, gene expression, *in vitro*, legumes, *Medicago truncatula*, water stress; *WEE1*

Introduction

Water stress can result in reducing crop yield world-wide (Boyer, 1982; Gonzalez *et al.*, 1995; Smirnov, 1993) and a recent UN survey has underlined the importance of water deficit in our planet and its effects for the coming generations unless urgent measures are taken. This situation is exacerbated in arid and semiarid ecosystems. Here legumes play a central agroecological role through their ability to use atmospheric nitrogen via the symbiosis with Rhizobia, and thus it reduces the need for fertilizers, improve food security, and generally favour the environment (Araújo *et al.*, 2015; Kohler *et al.*, 2008; Naya *et al.*, 2007; Rubio *et al.*, 2002; Ochatt, 2015). Studying a legume model species is thus timely and *Medicago truncatula* is of particular interest given its rather short life cycle and autogamy. It has a small and almost completely annotated genome (500–550 Mbp) which is publicly available (Goodstein *et al.*, 2012), and it is more drought tolerant than other legume crops such as pea, bean and soybean (Costa França *et al.*, 2000; Galvez *et al.*, 2005; Gonzalez *et al.*, 1998; Motan *et al.*, 1994). In spite of this, previous studies on water stress resistance in *M. truncatula* mostly concerned gene transfer (Alcântara *et al.*, 2015; Araújo *et al.*, 2015; Duque *et al.*, 2016). The assessment of physiological responses (Nunes *et al.*, 2008) and their genetic mechanisms (Badri *et al.*, 2011) is more limited.

Osmotic stress or water deficit can be defined as the absence of adequate moisture necessary for a plant to grow normally and complete its life cycle (Cabuslay *et al.*, 2002). Resistance mechanisms can be grouped into three categories: firstly escape, which enables the plant to complete its life cycle before the most intense period of water shortage, secondly avoidance, which prevents exposure to water stress, and finally tolerance, which enables the plant to withstand stress conditions (Golldack *et al.*, 2014; Levitt, 1972; Zhu, 2002). Some resistance mechanisms are constitutive and active before exposure to water shortage. In other cases, plants exposed to water stress alter their physiology, thereby acclimating themselves to withstand drier conditions. One of the tolerance mechanisms activated under such stress is that of mitigating osmotic stress, via the production of osmolytes such as proline, and soluble sugars, that protect cells against osmotic perturbation (Deinlein *et al.*, 2014; Choudhary *et al.*, 2005; Fulda *et al.*, 2011; Elmaghrabi *et al.*, 2013; Valliyodan and Nguyen, 2006). On the other hand, ϕ_w (water potential) is also known to induce a morphological variation in tissues subjected to osmotic stress, notably at the cellular level. Such variation is potentially useful to understand biodiversity by identifying cellular responses to stress that are not necessarily picked up by taxonomic or phylogenetic indices that consider cell shape or size *in vitro* (Ochatt *et al.*, 2008; Ochatt and Moessner, 2010). It is also important for assessing the competence for regeneration

in vitro (Ochatt *et al.*, 2008) following the recovery of tissues with a novel genetic makeup obtained via *in vitro* selection (Elmaghrabi *et al.*, 2013) or gene transfer (Alcântara *et al.*, 2015). Responses to abiotic stress factors involve a reprogramming of the expression of thousands of genes, which in turn result in the modification of a range of cellular and physiological processes (Cushman and Bohnert, 2000; Sreenivasulu *et al.*, 2004; Araújo *et al.*, 2015). One example of tolerance to stress at the molecular level, is the induction of *P5CS* that encodes Δ^1 -pyrroline - 5-carboxylate synthetase involved in proline biosynthesis (Silva-Ortega *et al.*, 2008). This gene is highly expressed in salt-and drought-tolerant plant species (Choudhary *et al.*, 2005) and it is induced under salt and water stress in many plant species including legumes (Chen *et al.*, 2009). The *P5CS* gene was also up-regulated in *M. truncatula* in response to salt stress (Elmaghrabi *et al.*, 2013). The kinetics of expression of genes involved in the cell cycle in plants exposed to high levels of abiotic stress has been the object of a number of studies (Gill and Tuteja, 2010; Roy, 2016; Zhao *et al.*, 2014). In Arabidopsis, a negative regulator of mitosis, *WEE1*, is strongly expressed in response to abiotic stress (De Schutter *et al.*, 2007). Osmotic stress imposed using PEG also up-regulated oxidative DNA damage and consequently DNA repair enzymes both in imbibed seeds (Balestrazzi *et al.*, 2011) and in plantlets (Macovei *et al.*, 2010). Our recent work with *M. truncatula* also showed an increased expression of *WEE1* and *CCS52* (*CELL CYCLE SWITCH PROTEIN 52*, another gene involved in the cell cycle) in salt-acclimated tissues as well as expression of genes involved in salt tolerance (*SOS1* encoding a Na^+/H^+ antiporter) and embryogenesis *in vitro* (*SOMATIC EMBRYOGENESIS RECEPTOR-LIKE KINASE 1*, *SERK1*) (Elmaghrabi *et al.*, 2013).

Tissue culture has been used in the selection of water stress tolerant cell lines that have been used to regenerate plants resistant to harsh environmental conditions in a range of crops including *Medicago sativa* L., tomato, soybean and wheat (Sakthivelu *et al.*, 2008; Guóth *et al.*, 2010; Mahmood *et al.*, 2012). Water deficit *in vitro* can be imposed through treatment with PEG 6000 (Ochatt *et al.*, 1998; Guóth *et al.*, 2010; Yang *et al.*, 2012; Rai *et al.*, 2011). The adsorbant property of this inert osmolyte provokes in plant cells and tissues the same or comparable effects to those obtained by drying soil at the same ϕ_w and without any other associated detrimental effects (Michel and Kaufmann, 1973). PEG 6000 thus closely mimics soil water stress (Lu *et al.*, 1998) and induces increases in total soluble sugars which serve as an osmoticum, or can be a source of respiratory substrates (Srivastava *et al.*, 1995; Elmaghrabi *et al.*, 2013). Additionally, PEG was shown to stimulate somatic embryogenesis *in vitro* (Attree *et al.*, 1995; Igasaki *et al.*, 2003). PEG 6000 was also used, and at similar concentrations as here

(although osmolarity was expressed in MPa rather than in mOsm/kg as in this work), in studies on PEG-induced DNA damage with *M. truncatula* *in vitro* plantlets (Macovei *et al.*, 2010) and seeds (Balestrazzi *et al.*, 2011).

As many legumes are grown (or have their center of origin) in regions with an arid to semi-arid climate (Smýkal *et al.*, 2015), a number of studies have identified genes, QTLs, ESTs and SNPs that are responsive to drought stress in several species (Jacob *et al.*, 2016). However, the molecular basis of water stress tolerance is not fully understood in *Medicago truncatula* (Araújo *et al.*, 2015). The aim of the current work was to examine the extent to which treatments with PEG could enhance osmotic stress tolerance potential in callus of *Medicago truncatula*. In addition, the accumulation of osmoprotectants, the effects on cell morphology (shape and size) and division competence, and the expression of *WEE1*, *CCSS52*, *P5CS* and *SOS1* were monitored in the PEG treatments to investigate the mechanism underlying the induced water stress responses as compared to those activated in response to salt stress (Elmaghrabi *et al.*, 2013). The expression of *SERK1* was also analyzed, given its key role on the competence for the subsequent regeneration through somatic embryogenesis of plants that may potentially carry the stress resistance trait acquired.

Materials and Methods

Plant material

Medicago truncatula cv. Jemalong line A17 ($2n = 2x = 16$, 1C value = 0.48 pg) was used in this study. One hundred leaves were explanted to tissue culture from 4 week old aseptically grown plants onto a medium consisting of MS basal medium (Murashige and Skoog, 1962) with 2 mg L⁻¹ NAA (alpha-naphthalene acetic acid; Sigma, Poole, UK) and 0.5 mg L⁻¹ BAP (6-benzylaminopurine; Sigma, Poole, UK), hereafter called MANA medium as in Elmaghrabi *et al.* (2013), and dispensed in multi-well dishes as 2 mL aliquots per well. Cultures were incubated at 24/22 °C (day/night), with a 16/8 h (light/ dark) photoperiod of 90 µmol m⁻² s⁻¹ from warm white fluorescent tubes. After 4 weeks, explants were sub-cultured on the medium above and the frequency of callus initiation assessed.

Leaf-derived embryogenic callus was obtained after culture on MANA medium for 5 months. Calli were screened for embryogenesis (i.e. somatic embryos at different developmental stages, identified as spherical glistening nodules when globular, through to elongated greening structures at later stages) or, organogenesis (development of shoots and/ or roots), as reported

elsewhere (Ochatt *et al.*, 1998; Elmaghrabi and Ochatt, 2006; Chen *et al.*, 2011; Ochatt *et al.*, 2013; Ochatt and Revilla, 2016). Only embryogenic calli were transferred onto 25 ml of MANA medium supplemented with or without 10 % w/v (-0.66 MPa) PEG6000 (PEG; Sigma, Poole, UK) for six months in order to acclimate the cultures under conditions that mimic water (osmotic) stress (at least 12 calli per treatment). This PEG concentration was chosen based on previous studies with various species (Biswas *et al.*, 2002, Guóth *et al.*, 2010) and also including *M. truncatula* (Macovei *et al.*, 2010, Balestrazzi *et al.*, 2011). Growth data (g fresh weight, g FW) were recorded and results were statistically analysed ($P \leq 0.05$; Kruskal Wallis followed by a Dunn's test).

Proline and water soluble carbohydrate measurements

Proline content was measured as described in Elmaghrabi *et al.* (2013) and according to Troll and Lindsley (1955) and Boukel and Houassine (1997) from callus tissue (100 mg per sample per treatment) grown on 0 and 10 % (w/v) PEG. All treatments were repeated three times. Optical density was measured using a spectrophotometer (UNICAM; Cambridge, UK) at a wavelength of 528 nm and calibrated using a standard curve of proline solutions (0.1–0.4 mg mL⁻¹; Sigma, Poole, UK).

Determination of soluble sugars was by the anthrone method (Elmaghrabi *et al.*, 2013; Plummer, 1987) using 100 mg callus samples from 0 and 10 % (w/v) PEG treatments (3 replicates). The soluble sugar content was measured spectrophotometrically (UNICAM, Cambridge, UK) at 585 nm and the data were converted to mg L⁻¹ using the calibrations established prior to the assay.

Medium and callus osmolarity

For measurements of medium and callus osmolarity a Wescor (model VAPRO 5520, South Logan, USA) vapour pressure micro-osmometer was used and a minimum of three 10 µL samples were measured. For medium osmolarity assessments, 10 mL of the medium were vortexed prior to collecting the 10µL samples to be measured. For callus osmolarity measurements, 1 g fresh weight of tissue was collected in 2 mL of liquid medium and centrifuged (100 g, 10 min, 10°C). The supernatant was carefully removed, the pellet was crushed in an Eppendorf with a pestle and centrifuged (1000g, 10 min, 4°C), and this second supernatant was finally employed for measurements of osmolarity. Results from such measurements, expressed in mMkg⁻¹, are the mean \pm S.E. of a minimum of three individual

samples per treatment, and were performed at the time of sub culturing and over at least three consecutive subcultures.

Mitotic index, cell viability and cell morphology

For determinations of C-value stability of calli following *in vitro* selection for several months they were compared to leaf tissues from the original plants. Nuclei were mechanically isolated from about 0.2 g of calli or from a single leaf of *M. truncatula* A17 grown in green house conditions. Tissues were chopped roughly with a sharp razor in 400µl of nuclei extraction buffer and 1.6 mL of staining buffer (Partec®; Canterbury, UK) (Ochatt, 2008). The suspension was filtered through a 20 µm nylon mesh and 4, 6 diamidino-2-phenylindole (DAPI; Sigma, Poole, UK), an A-T binding specific fluorochrome, was added to the filtrate to a final concentration of 1 µg mL⁻¹. The DNA contents of the isolated nuclei suspension were analysed using a Partec PAS-II flow cytometer equipped with an HBO-100 W mercury lamp and a dichroic mirror (TK420). Ten replicated calli for each treatment were analyzed, with a minimum of 3000 to 10000 nuclei per run. The mitotic index was calculated according to the formula: $MI = 4 \times 4C / \Sigma 2C + 4C$, where 2C and 4C correspond to the mean integrated value of nuclei in G1 phase and G2, respectively (Ochatt, 2008).

Cell viability was estimated by dual staining with fluorescein diacetate (FDA; Sigma, Poole, UK) and propidium iodide (PI; Sigma, Poole, UK). Cell suspensions (75 µL) from each treatment were mixed with 75 µL of dual staining solution containing FDA (200 µg mL⁻¹; Widholm, 1972) and propidium iodide (PI at 120 µg mL⁻¹) on ice and incubated for 20 min. The FDA molecule is cleaved by the esterases in the cytoplasm into acetate and fluorescein which, being hydrophilic accumulates in the cytoplasm of metabolically active (alive) cells that, upon excitation with the UV light fluoresce yellow-green, while dead cells appear red using a fluorescent microscope. A minimum of 300 cells are counted and results are expressed as the percentage of fluorescing cells referred to the total number of cells in the field.

For the cell morphology characterization, FDA stained slides of the control and PEG-treated cells were observed under the microscope under the UV. The surface area of cells and nuclei was determined at 2, 4 and 6 months of culture, using the image acquisition programmes ArchimedPlus and Histolab (Microvision, France) as reported (Ochatt *et al.*, 2008), and a shape coefficient (Ochatt and Moessner, 2010) was applied at 6 months of culture. Briefly, this shape coefficient (SC) is calculated based on the half length of the cell along its longest (a) and shortest (b) axes, as:

$$SC = \frac{\sqrt{a^2 - b^2}}{a}$$

For each treatment, nucleus and cell size were measured on 10 cells at 2 and 4 months of culture and at least 20 cells at 6 months of culture, and results expressed as the mean \pm SE.

The SC distinguishes round from elongated shapes, since SC values close to 1.0 correspond to elongated cells while SC values close to 0.5 correspond to rounder cell shapes.

Real time PCR

RNA was extracted and genomic DNA removed by DNase treatment (Elmaghrabi *et al.*, 2013; Spadafora *et al.*, 2012), and its absence verified using 18S rRNA primers (Spadafora *et al.*, 2011). Retrotranscription was carried out using an Ambion kit (RETROscript Reverse transcription for RT-PCR; Foster City, USA) and 2 μ g of RNA. An ABsoluteTM QPCR SYBR Green Mix (Thermo Scientific, Waltham, USA) kit was used for real time PCR. Reactions (in a total volume of 25 μ L) consisted of: 5 μ L cDNA (1:20 dilution), 12.5 μ L ABsoluteTM QPCR SYBR Green Mix, 1.75 μ L of each primer (10 μ M) and 4 μ L H₂O. Reactions were cycled in an MJ Research OPTICON 2 (Quebec, Canada), in triplicate under the following conditions: 95 °C for 10 min, 40 cycles of: 95 °C for 15 s, 60 °C for 30 s and 72 °C for 30 s and one cycle of 72 °C for 30 s. For testing primer specificity, a melting curve analysis was performed after amplification (from 60 to 98 °C with an increasing heat rate of 0.5 °C s⁻¹). A relative quantification of gene expression was calculated using the 2-DDCT method (Livak and Schmittgen, 2001). Primers for the target genes: *MtSOS1*, *MtWEE1*, *MtSERK1*, *MtP5CS* and *MtCCS52* are as described in Elmaghrabi *et al.* (2013). Mt18S rRNA primers were used to normalise the results as it was shown previously that 18S rRNA was a reliable reference gene for stress responses in *M. truncatula* (Elmaghrabi *et al.*, 2013), and widely used across a range of different species for developmental and stress-response studies (e.g. Price *et al.*, 2008; Wagstaff *et al.*, 2010).

Statistical Analyses

Unless otherwise stated above, data were analysed using R software (R version 3.3.2, Foundation for Statistical Computing). One or two way (as appropriate) ANOVA tests followed by a Tukey's test, or non-parametric statistical tests (Kruskal Wallis followed by a Dunn's test) were applied to determine differences across multiple samples. Comparisons between pairs of samples were performed using a Student's t-test, or if not normally distributed using a Wilcoxon signed rank test. Regression equation and R² value for the growth data were calculated in Excel.

Details of tests applied are provided in the legend to each Figure and all original data are provided as Supplementary materials (S1 to S6).

Results

PEG enhanced callus phenotype and embryogenic competence without reducing its viability

There were no significant differences between *Medicago truncatula* leaf callus cultured on MANA medium or MANA medium supplemented with 10% w/v PEG6000 over the 6 months of culture, and the linear growth rates for control and PEG treatments were 0.155 and 0.168 g month⁻¹ respectively (Figure 1). This only represents a 1.08 fold increase in the PEG compared with the control treatment indicating very similar rates of growth in each treatment irrespectively of the presence or absence of PEG in the culture medium. However, these data also suggest that callus tissues were PEG-tolerant already within 1 month of sub-culture and retained such tolerance throughout the experiment. Qualitative observations of callus indicated that those treated with PEG were typically bright green in colour and exhibited clear evidence of embryogenesis as did also the controls; however, the controls were brown in colour and the somatic embryos regenerated looked blocked at an early (globular to heart) developmental stage (Figure 2 A,B). The typical bright green coloration of calli in the PEG treatment would tend to indicate their robustness for both growth and embryogenesis regardless of the length of time in culture, and could also be ascribed to an increased tissue photochemistry linked to the tolerance acquired by onset of a priming process by the long-term culture on PEG. This was confirmed during the cell viability assessments with fluorescein diacetate where calli grown on 10% w/v PEG for 6 months contained 81.00 ± 1.5 % viable cells compared to 68.00 ± 5.2 for calli grown only on the MANA medium (Figure 2 C,D).

Mitotic index, cell and nuclear size increase, and cell shape changes in calli cultured on PEG-medium

Flow cytometry (FCM) was used to compare the C value distribution of cells from greenhouse grown *M. truncatula* leaves and callus derived from leaf material cultured for 6 months on 10% w/v PEG6000 (Figure 3). Flow cytometry raw profiles of leaves exhibited two peaks (Figure 3A), corresponding respectively to the nuclei in G1 phase (2C DNA) and those in G2/M (4C DNA), where their analysis after fitting them to Gauss curves resulted in a distribution of nuclei into three subpopulations as follows: G1 77.39%, S 17.32% and G2/M 5.29%, and coupled with a calculated mitotic index of 1.999 (Figure 3C). A very similar profile was obtained from calli cultured on MANA medium alone (not shown) which showed no obvious deviation from the

mother plant tissues from which they originated, while the flow cytometry profiles of calli cultured on PEG was very different (Figure 3B). In the 10% PEG6000 treatment, four peaks were typically detected consistent with 2, 4, 8 and 16 C populations (Figure 3B), and indicative of the occurrence of endoreduplication. The mitotic index was also significantly higher ($P \leq 0.05$) for the calli grown on PEG6000 containing medium (Figure 3C), which is also indicative of the onset of an endoreduplication phenomenon.

Interestingly, cell size also showed a significantly higher value for cells from calli grown on PEG6000 after just 2 months of culture (Figure 4A) and nuclear area was greater on PEG after 4 months (Figure 4B). This is consistent with the occurrence of endoreduplication, and this was coupled with a modified cell shape (Figure 4C), with cells grown on PEG6000 exhibiting a significantly lower SC than control cells. Thus, PEG-grown cells were consistently and significantly ($P < 0.05$) rounder ($SC = 0.608 \pm 0.117$) than control cells which were more elongated ($SC = 0.833 \pm 0.090$) (Figure 4D).

Osmolarity, proline and sugar levels rise following PEG treatment

After six months of 10% PEG 6000 treatment, there was a significant increase in osmolarity of callus in the PEG treatment compared with the control (Figure 5A) while osmolarity of the medium remained more constant. Proline and soluble sugar levels also increased significantly compared with the control (MANA without PEG) (Figure 5B,C). However there were no significant differences in water content between the PEG and control treatments (Figure 5D).

***MtWEE1* expression is highly up-regulated following PEG treatment while *MtSOS1* is down-regulated.**

We chose to examine the expression of five genes as markers of processes related to osmotic (water) stress. These comprised *MtSOS1* (salt stress response), *MtWEE1* (cell cycle checkpoint), *MtSERK1* (embryogenesis) *MtP5CS* (proline metabolism), and *MtCC52* (ploidy marker) in the embryogenic calli treated with PEG. Expression of these genes was measured using quantitative real time PCR after six months of callus culture in PEG6000 (10%) and compared with the control treatment (0 % PEG) and greenhouse grown leaves.

A highly significant reduction in the expression of *MtSOS1* occurred in the 10% PEG compared with the control treatment such that *MtSOS1* transcripts were virtually undetectable, comparable to expression in leaf (Figure 6A). Conversely, *MtWEE1* and *MtCC52* expression was significantly higher in the PEG treated calli compared with the control treatment which in turn

was higher than expression in leaf (Figure 6B,E). *MtSERK*, and *MtP5CS* were expressed significantly more in the control callus and 10% PEG treated callus but there was no significant difference in expression between the treated and untreated calli. (Figures 6C and D).

Discussion

Osmotic stress, provoked by insufficient ground and/or rain water, is a paramount constraint for plant growth and development. Cultures of callus on media that impose water deficit is a method for generating new, more tolerant, plants. Here we have shown that long term culture of *M. truncatula* calli on medium containing 10% PEG6000 to impose an osmotic stress results in the production of morphologically enhanced calli. An analysis of protective metabolite levels, cellular morphology, cell division and gene expression was undertaken to understand the effects of the imposed stress.

In this work, callus growth was not significantly different plus or minus 10% PEG over a period of six months, suggesting that calli on PEG acquired tolerance to osmotic (water) stress (Figure 1) probably mediated by an early osmotic adjustment which was likely associated to various modifications at the cellular level (Singh *et al.*, 2015). In fact, both cell viability and mitotic index were higher in the PEG treated cells compared to the control indicating a healthy and proliferating culture. It is likely that this sustained viability in the PEG treated cells is due to the activation of defence mechanisms that may include an activation of DNA repair as shown previously (Balestrazzi *et al.*, 2011). The similarity in growth rates following 1 month \pm PEG further stresses that tolerance was obtained relatively rapidly. This result differs from those of Biswas *et al.* (2002) who found that in rice, callus proliferation in the presence of PEG was greater than the controls in some genotypes, although this was at a much lower PEG concentration (5-15 gL⁻¹). However in two genotypes of wheat, one drought tolerant, the other drought sensitive, water deficit decreased only slightly in the sensitive compared with the tolerant genotype under water conditions and, it did not change significantly in either the sensitive or tolerant genotype \pm 400 mM PEG 6000 (100–400 mOsm; -0.976 MPa; Guóth *et al.*, 2010). Likewise in chili pepper cultures, where there was very good growth after 12 months in 5-10 % PEG8000 (0.57 MPa; Santos-Díaz and Ochoa-Alejo, 1994). Note that after six months of treatment, although osmolarity of the callus increased, osmolarity of the medium did not change since PEG is not metabolised. We decided to analyse both the medium and callus osmolarity as an indirect way of assessing the nutrient consumption from the medium by cells, which impacts their internal salt concentrations, as shown before with various species among which *M. truncatula* (Ochatt *et al.*, 2008). An increased cell osmolarity appeared also to be a

reliable early marker of embryogenic competence (Ochatt *et al.*, 2008; Elmaghrabi *et al.*, 2013). Thus the increased callus osmolarity and embryogenic capacity seen here are in line with previous observations.

In this work (Figure 4) PEG-induced stress resulted in a highly significant increase of the size of both nuclei and cells after 6 months of culture on selection medium. Remarkably, this was also coupled with a consistent and significant modification of the cell shape, reflected by the SC values observed, indicative of an increased elasticity of cell walls under PEG-induced osmotic stress. A similar modification of cell wall elasticity was observed in *M. truncatula* plants subjected to a severe drought stress (Nunes *et al.*, 2008) and in transgenic *M. truncatula* lines expressing the trehalose-6-Phosphate Synthase 1 (*AtTPS1*) from *Arabidopsis thaliana* with altered response to water deficit and recovery (Alcântara *et al.*, 2015). Taken together, these observations suggest a profound elastic modification of the cell walls of water stress tolerant cells, perhaps deriving from a modified ratio among cell wall fractions, and should be the object of future studies.

Similar levels of somatic embryogenesis were observed in the PEG and control calli, however, calli in the PEG treatment were distinctly green in colour compared with the control. This might be consistent with more robust embryogenic callus in the PEG compared to the control treatments (Figures 1 and 2), which is not surprising since MANA medium is not conducive to full maturation of the somatic embryos formed in *M. truncatula* (Ochatt *et al.*, 2013; Ochatt and Revilla, 2016). It may also reflect the fact that *M. truncatula* is adapted to semiarid conditions and even under severe drought stress, pigment content is not affected (Biswal, 1997; Nunes *et al.*, 2008). PEG also improved somatic embryogenesis in other species (Attree *et al.*, 1995; Igasaki *et al.*, 2003). Both control and PEG treatments resulted in similar levels of somatic embryogenesis, which was consistent with the similar expression levels of *MtSERK* in the two treatments. Note that *SERK1* is highly expressed during embryo induction and early somatic embryo development in *M. truncatula* (Nolan *et al.*, 2009) and in *Araucaria angustifolia* (Steiner *et al.*, 2012). The possible stimulation of somatic embryogenesis in response to PEG treatment is consistent with other reports showing stress-induction of this process (Karami and Saidi, 2010), which in *M. truncatula* may be linked to increases in ABA (Nolan and Rose, 1998).

The physiological, metabolic and gene expression responses of calli to PEG-induced osmotic stress mirrored those found under salt stress treatments (Elmaghrabi *et al.*, 2013) in some respects but not in others, as summarized in Figure 7. In contrast to NaCl treatment, PEG treatment did not result in any increase in water content of the calli compared with the control

although osmolarity did increase. This could be explained by the differential modes of action between NaCl and PEG, as the high MW of PEG exerts a constant osmotic pressure but does not allow its entry across the wall and hence avoids cell plasmolysis which results in different energy costs and different effects on growth (Munns, 2002). However soluble sugars did increase suggesting they are a useful marker of both salt and water (osmotic) stress (Figure 5). PEG also induced a high level of proline accumulation, which was far higher than the largest proline accumulation under stress induced by NaCl (Elmaghrabi *et al.*, 2013), indicating that this might also be a component of osmotic stress tolerance in *M. truncatula* (Figure 5B), as has been found in other species (Deinlein *et al.*, 2014). Validating this hypothesis would require field trials with regenerants from these cultures and goes beyond the scope of this study. However, the expression of *MtP5CS* in callus grown on PEG was similar to the control (Figure 6D) and hence does not correlate with increased levels of proline in the PEG treatment. This was surprising given that this gene encodes an enzyme that is central to proline synthesis and that its expression in *M. truncatula* cultures exposed to salt stress was elevated (Elmaghrabi *et al.*, 2013; Figure 7). It may suggest that this enzyme is not a key regulatory step in proline biosynthesis under these conditions, or that an initial rise in *MtP5CS* expression early in the culture period was sufficient to elevate proline concentrations and that after 6 months culture, increased gene expression was no longer necessary. In other words, whether modifying proline metabolism and the expression of genes involved in it, such as *P5CS*, may or not be used for engineering drought tolerance, and which approach should be adopted for such modification to be done remains uncertain (Bhaskara *et al.*, 2015).

More predictably, PEG did not induce *MtSOS1* expression which was down-regulated compared with the control. This gene is highly expressed in salt stress conditions as it encodes a protein that functions as a membrane-bound Na⁺ antiporter and contributes to Na⁺ depletion in the cytoplasm (Feki *et al.*, 2011; Smith *et al.*, 2010). Therefore the PEG data indicate a different (non ionic) pathway leading to osmotic stress tolerance compared with NaCl (ionic) tolerance (Figure 7) as reported by Elmaghrabi *et al.* (2013).

The expression of *MtCCS52* was upregulated by the PEG treatment (Figure 6E). In Arabidopsis this gene is a regulator of ploidy level and its expression is positively correlated to endoreduplication. In *M. truncatula* cultures exposed to long-term NaCl treatments this gene was up-regulated (Figure 7), alongside and increase in endoreduplication (Elmaghrabi *et al.*, 2013). Given the clear evidence for endoreduplication in the osmotic stress-resistant cultures here (Figure 3), the upregulation of *MtCCS52* is consistent with its role in Arabidopsis (Vanstraelen *et al.*, 2009).

That *MtWEE1* expression was more highly expressed in the PEG treatment suggests that this gene may have a role in maintaining normal growth in a treatment that mimics osmotic stress conditions. *WEE1* kinase might be necessary to regulate normal cell size in the face of ion toxicity and osmotic (non-ionic) water stress although this could only be resolved by exposing calli from *M. truncatula wee1* knockouts to these treatments. Alternatively, as a gene that is expressed in the DNA damage and DNA replication checkpoints, it may be induced in response to either single strand or double strand DNA breaks as it is in Arabidopsis (De Schutter *et al.*, 2007). However, Gonzalez *et al.* (2004) observed high expression of *LeWEE1* in tomato (*Lycopersicon esculentum* Mill.) which was correlated with endoreduplication during fruit development. Our results do not seem to indicate that there has been irreversible DNA damage due to the osmotic stress imposed on callus, and it would therefore be legitimate to link this to *WEE1* expression and its role in replication checkpoint and DNA damage and the possibility that the PEG concentration used and the long-term culture on it resulted in priming (Singh *et al.*, 2015). Other genes could also been involved in the process though, and, in this respect, in order to protect their gene integrity from DNA damage plants are capable of activating a specific response system that regulates the cell cycle, but also DNA repair and programmed cell death where genes such as *Suppressor Of Gamma response 1 (SOG1)* (Yoshiyama *et al.*, 2014) and *Breast Cancer 1 (BRCA1)* (Block-Schmidt *et al.*, 2011) are known to play a central role in DNA repair, chromosome segregation and chromatin remodeling So the increase in *MtWEE1* seen here may be both linked to the increase in endoreduplication and required to protect the cells from DNA damage induced by the PEG-induced osmotic stress treatment.

Conclusion

The data reported in this study of responses to 10% PEG compared with no PEG controls indicates that at this level of osmotic stress it is possible to induce a high level of embryogenesis with no penalty on growth rate. This appears to be achieved by up-regulating protective mechanisms such as the production of osmoprotectant solutes and switching on the expression of *MtWEE1*. The increase in *MtWEE1* and *MtCCS52* expression may cause an increase in endoreduplication while protecting the cells against the potentially damaging effects of the osmotic stress on DNA integrity.

Acknowledgments

AME thanks the Libyan Government for financial support, and the authors thank Mike O'Reilly for assistance with analytical work.

References

- Alcântara, A., Morgado, R.S., Silvestre, S., Marques da Silva, J., Bernardes da Silva, A., Fevereiro, P., Araújo, S.S. (2015) A method to identify early-stage transgenic *Medicago truncatula* with improved physiological response to water deficit. *Plant Cell Tiss. Organ Cult.* 122:605–616. <http://dx.doi.org/10.1007/s11240-015-0793-4>
- Araújo, S.S., Beebe, S., Crespi, M., Delbreil, B., González, E.M., Gruber, V., Lejeune-Henaut, I., Link, W., Monteros, M.J., Prats, E., Rao, I., Vadez, V., Vaz Patto, M.C. (2015) Abiotic stress responses in legumes: strategies used to cope with environmental challenges. *Crit. Rev. Plant Sci.* 34:237–280. <http://dx.doi.org/10.1080/07352689.2014.898450>
- Attree, S.M., Pomeroy, M.K., Fowke, L.C. (1995) Development of white spruce (*Picea glauca* (Moench.) Voss) somatic embryos during culture with abscisic acid and osmoticum, and their tolerance to drying and frozen storage. *J. Exp. Bot.* 46: 433–439. <https://doi.org/10.1093/jxb/46.4.433>
- Badri, M., Chardon, F., Huguet, T., and Aouani, M. E. (2011) Quantitative trait loci associated with drought tolerance in the model legume *Medicago truncatula*. *Euphytica* 181: 415–428. Doi:10.1007/s10681-011-0473-3
- Balestrazzi A, Confalonieri M, Macovei A, Carbonera D (2011) Seed imbibition in *Medicago truncatula* Gaertn.: Expression profiles of DNA repair genes in relation to PEG-mediated stress. *Journal of Plant Physiology* 168 706–713. doi:10.1016/j.jplph.2010.10.008
- Bhaskara, G.B., Yang, T.-H., Verslues, P.E. (2015) Dynamic proline metabolism: importance and regulation in water limited environments. *Front Plant Sci.* 2015; 6: 484. doi: 10.3389/fpls.2015.00484
- Biswal B. Chloroplast metabolism during leaf greening and degreening. In: Pessarakli M, editor. *Handbook of photosynthesis*. New York- Basel-Hong Kong: Marcel Dekker; 1997. pp. 72–81.
- Biswas, J., Chowdhury, B., Bhattacharya, A., Mandal, A.B. (2002) In vitro screening for increased drought tolerance in rice. *In Vitro Cell Dev Biol -Plant* 38: 525-530. DOI : 10.1079/IVP2002342

Block-Schmidt AS, Dukowic-Schulze S, Wanieck K, Reidt W, Puchta H (2011) BRCC36A is epistatic to BRCA1 in DNA crosslink repair and homologous recombination in *Arabidopsis thaliana*. *Nucleic Acids Research* 39:146–154. doi:10.1093/nar/gkq722

Boukel, M., Houassine, D. (1997) Adaptation au stress hydrique de quelques variétés de blé dur (*Triticum durum*). Thèse magistère, INA, Algérie: 90.

Boyer, J.S. (1982) Plant productivity and environment. *Science*. 218: 443-448. DOI: 10.1126/science.218.4571.443

Cabuslay, S.G., Ito, O., Alejar, A.A. (2002) Physiological evaluation of responses of rice (*Oryza sativa* L.) to water deficit. *Plant Sci*. 163: 815-827. [https://doi.org/10.1016/S0168-9452\(02\)00217-0](https://doi.org/10.1016/S0168-9452(02)00217-0)

Chen, J.-B., Wang, S.-M., Jing, R.-L., Mao, X.-G. (2009) Cloning the PvP5CS gene from common bean (*Phaseolus vulgaris*) and its expression patterns under abiotic stresses. *J. Plant Physiol*. 166:12–19. doi: 10.1016/j.jplph.2008.02.010

Choudhary, N.L., Sairam, R.K., Tyagi, A. (2005) Expression of Δ^1 -pyrroline -5 carboxylate synthetase gene during drought in rice (*Oryza sativa* L.). *Indian J. Biotech. Biophys.* 42: 366-370.

Costa França, M.G., Thi, A.T.P., Pimentel, C., Rossiello, R.O.P., Zuily-Fodil, Y., Laffray, D. (2000) Differences in growth and water relations among *Phaseolus vulgaris* cultivars in response to induced drought stress. *Environ. Exp. Bot.* 43:227–237. doi: 10.1016/S0098-8472(99)00060-X

Cushman, J.C., Bohnert, H.J. (2000) Genomic approaches to plant stress tolerance. *Curr. Opin. Plant Biol.* 3: 117-124. DOI: 10.1016/S1369-5266(99)00052-7

Deinlein, U., Stephan, A.B., Horie, T., Luo, W., Xu, G., Schroeder, J.I. (2014) Plant salt-tolerance mechanisms *Tr. Plant Sci.* 19: 371–379. doi: 10.1016/j.tplants.2014.02.001

De Schutter, K., Joubes, J., Cools, T., Verkest, A., Corellou, F., Babiychuk, E., Der Schueren, E.-V., Beeckman, T., Kushnir, S., Inze, D., De Veylder, L. (2007) *Arabidopsis* WEE1 Kinase Controls Cell Cycle Arrest in Response to Activation of the DNA Integrity Checkpoint. *The Plant Cell*. 19: 211–225. doi:10.1105/tpc.106.045047

Duque, A.S., López-Gómez, M., Kráčmarová, J., Gomes, C.N., Araújo, S.S., Lluch, C., Fevereiro, P. (2016) Genetic engineering of polyamine metabolism changes *Medicago*

497 *truncatula* responses to water deficit. Plant Cell Tiss. Organ Cult. 127: 681-690.
 498 <https://doi.org/10.1007/s11240-016-1107-1>

499 Elmaghrabi, A.M., Ochatt, S.J. (2006) Isoenzymes and flow cytometry for the assessment of
 500 true-to-typeness of calluses and cell suspension of barrel medic prior to regeneration. Plant
 501 Cell Tiss. Organ Cult. 85: 31-43. doi:10.1007/s11240-005-9046-2

502 Elmaghrabi, A.M., Ochatt, S.J., Rogers, H., Francis, D. (2013) Enhanced tolerance to salinity
 503 following cellular acclimation to increasing NaCl levels in *Medicago truncatula*. Plant Cell
 504 Tiss. Organ Cult. 114: 61-70. doi:10.1007/s11240-013-0306-2

505 Feki, k., Quintero, F.J., Pardo, J.M., Masmoudi, K. (2011) Regulation of durum wheat Na⁺/H⁺
 506 exchanger TdSOS1 by phosphorylation. Plant Mol Biol. 76:545–556. doi:10.1007/s11103-
 507 011-9787-8.

508 Fulda, S., Mikkat, S., Stegmann, H., Horn, R. (2011) Physiology and proteomics of drought
 509 stress acclimation in sunflower (*Helianthus annuus* L.). Plant Biol. 13: 632-642. doi:
 510 10.1111/j.1438-8677.2010.00426.x.

511 Galvez, L., Gonzalez, E.M., Arrese-Igor, C. (2005) Evidence for carbon flux shortage and
 512 strong carbon/nitrogen interaction in pea nodules at early stages of water stress. J Exp. Bot.
 513 56: 2551-2561. <https://doi.org/10.1093/jxb/eri249>

514 Gill, S.S., Tuteja, N. (2010) Reactive oxygen species and antioxidant machinery in abiotic stress
 515 tolerance in crop plants. Plant Physiol. Biochem. 48: 909-930. doi:
 516 10.1016/j.plaphy.2010.08.016

517 Golldack, D., Li, C., Mohan, H., Probst N. (2014) Tolerance to drought and salt stress in plants:
 518 unraveling the signaling networks Front. Pl. Sci. <http://dx.doi.org/10.3389/fpls.2014.00151>

519 Gonzalez, E.M., Aparicio-Tejo, P.M., Gondon, A.J., Minchin, F.R., Royuela, M., Arrese-Igor,
 520 C. (1998) Water-deficit effects on carbon and nitrogen metabolism of pea nodules. J Exp. Bot.
 521 49: 1705-1714. <https://doi.org/10.1093/jxb/49.327.1705>

522 Gonzalez, E.M., Cordon., A.J., James, C.L., Arrese-Igor., C. (1995) The role of sucrose
 523 synthase in the response of soybean nodules to drought. J Exp. Bot. 46: 1515-1523.
 524 <https://doi.org/10.1093/jxb/46.10.1515>

525 Gonzalez, N., Hernould, M., Delmas, F., Gevaudant, F., Duffe, P., Causse, M., Mouras, A.,
 526 Chevalier, C. (2004) Molecular characterization of *WEE1* gene homologue in tomato

527 (*Lycopersicon esculentum* Mill). Plant Mol. Biol. 56: 849-861. doi:10.1007/s11103-004-
528 5110-2

529 Goodstein, D.M., Shu, S., Howson, R., Neupane, R., Hayes, R.D., Fazo, J., Mitros, T., Dirks,
530 W., Hellsten, U., Putnam, N., Rokhsar, D.S. (2012) Phytozome: a comparative platform for
531 green plant genomics. Nucleic Acids Res 40:D1178–D1186. doi: 10.1093/nar/gkr944

532 Guóth, A., Benyo, D., Csiszar, J., Galle, A., Horvath, F., Cseuz, L., Erdei, L., Tari, I. (2010)
533 Relationship between osmotic stress-induced abscisic acid accumulation, biomass production
534 and plant growth in drought-tolerant and -sensitive wheat cultivars. Acta Physiol. Plant. 32:
535 719–727. doi:10.1007/s11738-009-0453-6

536 Igasaki, T., Sato, T., Akashi, N., Mohri, T., Maruyama, E., Kinoshita, I., Walter, C., Shinohara,
537 K. (2003) Somatic embryogenesis and plant regeneration from immature zygotic embryos of
538 *Cryptomeria japonica* D. Don. Plant Cell Rep. 22: 239-243. doi:10.1007/s00299-003-0687-
539 5

540 Jacob, C., Carrasco, B., Schwember, A.R. (2016) Advances in breeding and biotechnology of
541 legume crops. Plant Cell Tiss. Organ Cult. 127: 561-584. doi:10.1007/s11240-016-1106-2

542 Karami, O., Saidi, A. (2010) The molecular basis for stress-induced acquisition of somatic
543 embryogenesis. Molecular Biology Reports 37, 2493–2507. doi: 10.1007/s11033-009-9764-
544 3.

545 Kohler J, Hernández JA, Carabaca F, Roldán A. (2008) Plant growth promoting rhizobacteria
546 and arbuscular mycorrhizal fungi modify alleviation biochemical mechanism in water stressed
547 plants. Funct Plant Biol. 35:141–151. doi: 10.1071/FP07218.

548 Levitt, J. (1972) Responses of plants to environmental stresses. Academic Press, New York.
549 ISBN : 0124455018

550 Livak KJ, Schmittgen TD (2001) Analysis of relative gene expression data using real-time
551 quantitative PCR and the 2-DDC method. Methods 25:402–408. Doi:
552 10.1006/meth.2001.1262

553 Lu, Z., Neumann, P.M. (1998) Water-stressed maize, barley and rice seedlings show species
554 diversity in mechanisms of leaf growth inhibition. J. Exp. Bot. 49: 1945-1952.
555 <https://doi.org/10.1093/jxb/49.329.1945>

Macovei A, Balestrazzi A, Confalonieri M, Carbonera D. (2010) The Tdp1 (Tyrosyl-DNA phosphodiesterase) gene family in barrel medic (*Medicago truncatula* Gaertn.): bioinformatic investigation and expression profiles in response to copper- and PEG-mediated stress. *Planta* 232:393–407. Doi: 10.1007/s00425-010-1179-9

Mahmood, I., Razzaq, A., Hafiz, A.I., Kaleem, S., Khan, A.A., Qayyum, A., Ahmad, M. (2012) Interaction of callus selection media and stress duration for *in vitro* selection of drought tolerant callus of wheat. *Afr. J. Biotech.* 11: 4000-4006. DOI: 10.5897/AJB11.2610

Michel, B.E., Kaufmann, M.R. (1973) The osmotic potential of polyethylene glycol 6000. *Plant Physiol.* 51: 914–916. DOI: <https://doi.org/10.1104/pp.51.5.914>

Motan, J.F., Becana, M., Iturbeormaetxe, I., Frechilla, S., Klucas, R.V., Aparicio Tejo, P. (1994) Drought induces oxidative stress in pea-plants. *Planta.* 194: 346-352. <https://doi.org/10.1007/BF00197534>

Munns R. (2002) Comparative physiology of salt and water stress. *Plant Cell Environ* 25:239–50. <http://dx.doi.org/10.1046/j.0016-8025.2001.00808.x>

Murashige, T., Skoog, F. (1962) A revised medium for rapid growth and bioassays with tobacco tissue cultures. *Physiol. Plant.* 15: 473-497. doi/10.1111/j.1399-3054.1962

Naya, L., Ladrera, R., Ramos, J., Gonzalez, E.M., Arrese-Igor, C., Minchin, F.R., Becana, M. (2007) the response of carbon metabolism and antioxidant defenses of alfalfa nodules to drought stress and to the subsequent recovery of plants. *Plant Physiol.* 144: 1104-1114. doi: 10.1104/pp.107.099648

Nolan, K.E., Rose, R.J. (1998) Plant regeneration from cultured *Medicago truncatula* with particular reference to abscisic acid and light treatments. *Aust. J. Bot.* 46:151–160. <https://doi.org/10.1071/BT96138>

Nolan, K.E., Kurdyukov, S., Rose, R.J. (2009) Expression of the somatic embryogenesis receptor like kinase1 (SERK1) gene is associated with developmental change in the life cycle of model legume *Medicago truncatula*. *J Exp. Bot.* 60:1759–1771. <https://doi.org/10.1093/jxb/erp046>

Nunes, C.M.J., Araújo, S.S., Marques da Silva, J., Fevereiro, P., Bernandes da Silva, A.R. (2008) Physiological responses of the legume model *Medicago truncatula* cv. Jemalong to water deficit. *Environ. Exp. Bot.* 63 : 289–296. <https://doi.org/10.1016/j.envexpbot.2007.11.004>

587 Ochatt, S.J., Marconi, P.L., Radice, S., Arnozis, P.A., Caso, O.H. (1998) *In vitro* recurrent
588 selection of potato: production and characterization of salt tolerant cell lines and plants. Plant
589 Cell Tiss. Org. Cult. 55: 1-8. <https://doi.org/10.1023/A:1026426331722>

590 Ochatt, S.J. (2008) Flow cytometry in plant breeding. Cytometry A 73:581-98.
591 doi:10.1002/cyto.a.20562

592 Ochatt, S., Muilu, R., Ribalta, F. (2008) Cell morphometry and osmolarity as early indicators
593 of the onset of embryogenesis from cell suspension cultures of grain legumes and model
594 systems. Plant Biosyst. 142:480-486. doi:10.1080/11263500802410827

595 Ochatt, S.J., Moessner, A. (2010) Rounding up plant cells. Int J Plant Biol 1:e8.
596 doi:10.4081/pb.2010.e8.

597 Ochatt, S., Jacas, L., Patat-Ochatt, E.M., Djennane, S. (2013) Flow cytometric analysis and
598 molecular characterization of *Agrobacterium tumefaciens*-mediated transformants of
599 *Medicago truncatula*. Plant Cell Tissue Organ Cult. 113: 237-244. doi:10.1007/s11240-012-
600 0263-1

601 Ochatt, S.J. (2015) Agroecological impact of an in vitro biotechnology approach of embryo
602 development and seed filling in legumes. Agron Sustain Dev 35:535–552. DOI
603 10.1007/s13593-014-0276-8

604 Ochatt, S.J., Revilla M.A. (2016) From Stress to Embryos: Some of the Problems for Induction
605 and Maturation of Somatic Embryos. In: In Vitro Embryogenesis in Higher Plants. Eds M.A.
606 Germanà, M. Lambardi. Methods in Molecular Biology 1359. Springer Protocols. Humana
607 Press, New York, Heidelberg, Dordrecht, pp. 523-536. DOI 10.1007/978-1-4939-3061-6

608 Plummer, D.T. (1987) Introduction aux techniques de biochimie, Ed.McGraw-Hill. Paris: 331.

609 Price, A., Orellana, D., Salleh, F., Stevens, R., Acock, R., Buchanan-Wollaston, V., Stead, A.
610 and Rogers, H.J. (2008) A comparison of leaf and petal senescence in wallflower reveals
611 common and distinct patterns of gene expression and physiology, Plant Physiol., 147: 1898-
612 1912. doi: 10.1104/pp.108.120402

613 Rai M. K., Kaliaa R. K., Singh R., Gangola M. P., Dhawan A.K. (2011) Developing stress
614 tolerant plants through in vitro selection—An overview of the recent progress. Environ. Exp.
615 Bot. 71: 89–98. doi: 10.1016/j.envexpbot.2010.10.021

616 Roy, S. (2016) Function of MYB domain transcription factors in abiotic stress and epigenetic
 617 control of stress response in plant genome. *Plant Signaling & Behavior* 11: e1117723.
 618 <http://doi.org/10.1080/15592324.2015.1117723>

619 Rubio, M.C., González, E.M., Minchin, F.R., Webb, K.J., Arrese-Igor, C., Ramos, J., Becana,
 620 M. (2002) Effects of water stress on antioxidant enzymes of leaves and nodules of transgenic
 621 alfalfa overexpressing superoxide dismutases. *Physiol. Plant* 115: 531-540. DOI:
 622 10.1034/j.1399-3054.2002.1150407.x

623 Sakthivelu, G., Akitha Devi, M.K., Giridhar, P., Rajasekaran, T., Ravishankar, G.A., Nedev,
 624 T., Kosturkova, G. (2008) Drought-induced alterations in growth, osmotic potential and *in*
 625 *vitro* regeneration of soybean cultivars. *Gen. Appl. Plant Physiol.* 34: 103-112.

626 Santos-Díaz, M. & Ochoa-Alejo, N. (1994) PEG-tolerant cell clones of chili pepper: Growth,
 627 osmotic potentials and solute accumulation. *Plant Cell, Tissue and Organ Culture* 37: 1-8.
 628 <https://doi.org/10.1007/BF00048110>

629 Silva-Ortega, C.O., Ochoa-Alfaro, A.E., Reyes-Agüero, J.A., Aguado-Santacruz, G.A.,
 630 Jiménez-Bremont, J.F. (2008) Salt stress increases the expression of p5cs gene and induces
 631 proline accumulation in cactus pear. *Plant Physiol. Biochem.* 46:82–92. DOI:
 632 10.1016/j.plaphy.2007.10.011

633 Singh, N. B., Singh, D., & Singh, A. (2015). Biological seed priming mitigates the effects of
 634 water stress in sunflower seedlings. *Physiology and Molecular Biology of Plants*, 21(2), 207–
 635 214. <http://doi.org/10.1007/s12298-015-0291-5>

636 Smirnoff N. The role of active oxygen in the response of plants to water deficit and desiccation.
 637 *New Phytol.* 1993;125:27–58. doi: 10.1111/j.1469-8137.1993.tb03863.x.

638 Smith, M.A., Coupland, G., Dolan, L., Harberd, N., Jones, J., Martin, C., Sablowski, R., Amey,
 639 A. (2010) *Plant biology (Eds) Environmental stress, Salt stress*. Published by Garland Science.
 640 Taylor & Francis Group. LLC. New York. USA: 472-479.

641 Smýkal, P., Coyne, C.J., Ambrose, M.J., Maxted, N., Schaefer, H., Blair, M.W., Berger, J.,
 642 Greene, S.L., Nelson, M.N., Besharat, N., Vymyslický, T., Toker, C., Saxena, R.K.,
 643 Roorkiwal, M., Pandey, M.K., Hu, J., Li, Y.H., Wang, L.X., Guo, Y., Qiu, L.J., Redden, R.J.,
 644 Varshney, R.K. (2015) Legume crops phylogeny and genetic diversity for science and
 645 breeding. *Crit. Rev. Pl. Sci.* 34:43–104. doi:[10.1080/07352689.2014.897904](http://doi.org/10.1080/07352689.2014.897904)

646 Spadafora, N.D., Doonan, J.H., Herbert, R.J., Bitonti, N.B., Wallace, E., Rogers, H.J., Francis,
647 D. (2011) Arabidopsis T-DNA insertional lines for CDC25 are hypersensitive to hydroxyurea
648 but not to zeocin or salt stress. *Ann. Bot.* 107:1183–1192. doi: 10.1093/aob/mcq142

649 Spadafora, N.D., Parfitt, D., Li, S., Bruno, L., Vaughan, R., Buchanan-Wollaston, V., Herbert,
650 R.J., Bitonti, M.B., Doonan, J., Albani, D., Prinsen, E., Francis, D., Rogers, H.J. (2012)
651 Perturbation of cytokinin and ethylene-signalling pathways explain the strong rooting
652 phenotype exhibited by Arabidopsis expressing the *Schizosaccharomyces pombe* mitotic
653 inducer, cdc25. *BMC Plant Biol.* 12:45. doi:10.1186/1471-2229-12-45

654 Sreenivasulu, N., Varshney, R.K., Kavi Kishor, P.B., and Weschke, W. (2004) Tolerance to
655 abiotic stress in cereals: a functional genomics approach. In: Gupta, P.K., Varshney, R.K.
656 (Eds), *Cereal Genomic*. 483-514. Doi: 10.1007/1-4020-2359-6_16

657 Srivastava, D. K., Gupta, V. K., Sharma, D. R. (1995) *In vitro* selection and characterization of
658 water stress tolerant callus cultures of tomato (*Lycopersicon esculentum* L.). *Ind. J. Plant*
659 *Physiol.* 38: 99–104.

660 Steiner, N., Santa-Catarina, C., Guerra, M.P., Cutri L., Dornelas, M.C., Floh, E.I.S. (2012) A
661 gymnosperm homolog of *SOMATIC EMBRYOGENESIS RECEPTOR-LIKE KINASE-1*
662 (*SERK1*) is expressed during somatic embryogenesis. *Plant Cell Tiss Organ Cult* 109: 41-50.
663 <https://doi.org/10.1007/s11240-011-0071-z>

664 Troll, W., Lindsley, J., (1955) A photometric method for the determination of proline. *J. Biol.*
665 *Chem.* 215: 655-660. <http://www.jbc.org/content/215/2/655.citation>

666 Valliyodan, B., Nguyen, H.T. (2006) Understanding regulatory networks and engineering for
667 enhanced drought tolerance in plants. *Curr. Opin. Plant Biol.* 9: 189-195. DOI:
668 10.1016/j.pbi.2006.01.019

669 Vanstraelen, M., Baloban, M., Da Ines, O., Cultrone, A., Lammens, T., Boudolf, V., Brown,
670 S.C., De Veylder, L., Mergaert, P. and Kondorosi E. (2009) APC/CCCS52A complexes
671 control meristem maintenance in the *Arabidopsis* root. *Proc Nat Acad Sci USA* 106: 11806-
672 11811. doi: 10.1073/pnas.0901193106

673 Wagstaff, C., Bramke, I., Breeze, E., Thornber, S., Harrison, E., Thomas, B., Buchanan-
674 Wollaston, V., Stead, T. and Rogers, H. (2010) A specific group of genes respond to cold
675 dehydration stress in cut *Alstroemeria* flowers whereas ambient dehydration stress accelerates

developmental senescence expression patterns, J. Exp. Bot., 61: 2905-2921.
<https://doi.org/10.1093/jxb/erq113>

Widholm, J.M. (1972) The use of fluorescein diacetate and phenosaphranine for determining the viability of cultured cells. Stain Technol. 47: 189-194.
[doi/abs/10.1093/stn/47.3.189](https://doi.org/10.1093/stn/47.3.189)

Yang, L., Li, Y., Shen, H. (2012) Somatic embryogenesis and plant regeneration from immature zygotic embryo cultures of mountain ash (*Sorbus pohuashanensis*). Plant Cell. Tiss. Organ Cult. 109 : 547-556. <https://doi.org/10.1007/s11240-012-0121-1>

Yoshiyama, K.O., Kimura, S., Maki, H., Britt, A.B., Umeda, M. (2014) The role of SOG1, a plant-specific transcriptional regulator, in the DNA damage response. Plant Sign & Behavior 9: e28889. PMID: 24736489; <http://dx.doi.org/10.4161/psb.28889>

Zhao, L., Wang, P., Hou, H., Zhang, H., Wang, Y., Yan, S., Huang, Y., Li, H., Tan, J., Hu, A., Gao, F., Zhang, Q., Li, Y., Zhou, H., Zhang, W., Li, L. (2014) Transcriptional Regulation of Cell Cycle Genes in Response to Abiotic Stresses Correlates with Dynamic Changes in Histone Modifications in Maize. PLoS ONE 9(8): e106070.
[doi:10.1371/journal.pone.0106070](https://doi.org/10.1371/journal.pone.0106070)

Zhu, J.-K. (2002) Salt and drought stress signal transduction in plants. Ann Rev Plant Biol 53:2273–2467. DOI: 10.1146/annurev.arplant.53.091401.143329

Figure Legends

Figure 1. Mean (\pm S.D.) callus growth over 6 months on MANA medium with 10% w/v PEG compared with control (MANA medium). Different letter combinations indicate significant differences ($P \leq 0.05$; Kruskal Wallis followed by a Dunn's test) ($n > 8$). See Supplementary File S1 for data and statistical analyses.

Figure 2. Callus phenotype and viability after 6 months on 10% w/v PEG and control media: (A) control callus; (B) PEG-selected callus, arrows indicate somatic embryos; viability of control callus after FDA staining observed under transmission (C) or UV (D) light; viability of PEG-selected callus after FDA staining observed under transmission (E) or UV (F) light. Scale bars are A = 2.13 mm, B = 3.34 mm; and C through to F = 200 μ m.

Figure 3. FCM profile and % distribution of nuclei in G1 (1st peak), G2 (2nd peak), S phase (trough between 1st and 2nd peaks) and polyploid nuclei (3rd (8C) and 4th (16C) peaks) from (A) leaves of *Medicago truncatula*, (B) *M. truncatula* callus cultured on 10% w/v PEG6000 for 6 months; (C) Mean (\pm S.D.) mitotic index in leaves compared to callus tissues after 9 continuous months growth on 10% w/v PEG. Different letters indicate significant differences ($P \leq 0.05$); $n=10$. See Supplementary File S2 for data and statistical analyses.

Figure 4. Effects of PEG6000 on cell morphology. (A) cell and (B) nuclear size (μ m²) at different time points during *in vitro* selection for PEG6000 (10%) resistance. Data are the means \pm S.D. from $n = 9$ replicates at 2 and 4 months of culture and $n = 22$ measurements at 6 months of culture. Bars with different letters were significantly different (Kruskal Wallis followed by Dunn's test) at different time points across treatments ($P < 0.05$). See Supplementary File S3 for data and statistical analyses. (C) Shape coefficient (SC) of cells from control (blue) and PEG-grown (orange) calli ($n = 22$) at 6 months of mean \pm S. D. (* indicates $P < 0.05$, Welch Two Sample t-test). See Supplementary File S4 for data and statistical analyses. (D) Images of cells from PEG and control cultures at 6 months of culture (scale bars PEG = 100 μ m, Control 200 μ m)

Figure 5. Comparisons between control and 10% PEG treatments after six months *in vitro* culture in terms of: (A) medium and callus osmolarity, (B) proline content, (C) soluble sugars, and (D) water content. Note different letters indicate significant differences ($P \leq 0.05$) between

treatments ($n=3 \pm \text{SD}$); ns = non-significant. (Kruskal Wallis followed by Dunn's test (A); Wilcoxon signed rank test (B) and Welch Two Sample t-test (C and D)). See Supplementary File S5 for data and statistical analyses.

Figure 6. Gene expression after 6 months culture on 10% PEG medium and control (MANA medium) with leaf as reference: (A) *MtSOS1*, (B) *MtWEE1*, (C) *MtSERK*, (D) *MtP5CS* and (E) *MtCCS52*. Different letters indicate significant differences amongst treatments/tissues ($P \leq 0.05$; 1-way ANOVA; $n = 3 \pm \text{SD}$).

Figure 7. Summarized effects on various parameters of imposing abiotic stress on embryogenic callus of *Medicago truncatula* following *in vitro* selection. Control callus tissues are compared with NaCl-tolerant callus (Elmaghrabi et al. 2013) and PEG-induced osmotic stress tolerant callus tissues in this study. Blue color indicates increase/upregulation. Red color indicates decrease/downregulation. When non-significant compared to controls white is used.

Supplementary material: Data and statistical analyses

S1: Data for callus growth of control and PEG treatments shown in Figure 1

S2: Data for Mitotic Index of control and PEG treatments shown in Figure 3(C)

S3 - Data for cell and nuclear area of control and PEG treatments shown in Figure 4(A) and 4(B)

S4 - Data for Shape Coefficient calculations of control and PEG treatments shown in Figure 4(C)

S5 - Data for control and PEG treatments shown in Figure 5: (A) osmolarity of media and callus; (B) proline accumulation; (C) soluble sugars; (D) water content

S6 -Data for Realtime PCR of control and PEG treatments shown in Figure 6




Figure 1. Mean (\pm S.D.) callus growth over 6 months on MANA medium with 10% w/v PEG compared with control (MANA medium). Different letter combinations indicate significant differences ($P \leq 0.05$; Kruskal Wallis followed by a Dunn's test) ($n > 8$). See Supplementary File S1 for data and statistical analyses.




Figure 2. Callus phenotype and viability after 6 months on 10% w/v PEG and control media: **(A)** control callus; **(B)** PEG-selected callus, arrows indicate somatic embryos; viability of control callus after FDA staining observed under transmission **(C)** or UV **(D)** light; viability of PEG-selected callus after FDA staining observed under transmission **(E)** or UV **(F)** light. Scale bars are A = 2.13 mm, B = 3.34 mm; and C through to F = 200 μm.




Figure 3. FCM profile and % distribution of nuclei in G1 (1st peak), G2 (2nd peak), S phase (trough between 1st and 2nd peaks) and polyploid nuclei (3rd (8C) and 4th (16C) peaks) from (A) leaves of *Medicago truncatula*, (B) *M. truncatula* callus cultured on 10% w/v PEG6000 for 6 months; (C) Mean (\pm S.D.) mitotic index in leaves compared to callus tissues after 9 continuous months growth on 10% w/v PEG. Different letters indicate significant differences ($P \leq 0.05$); $n=10$. See Supplementary File S2 for data and statistical analyses.




Figure 4. Effects of PEG6000 on cell morphology. **(A)** cell and **(B)** nuclear size (μm^2) at different time points during *in vitro* selection for PEG6000 (10%) resistance. Data are the means \pm S.D. from $n = 9$ replicates at 2 and 4 months of culture and $n = 22$ measurements at 6 months of culture. Bars with different letters were significantly different (Kruskal Wallis followed by Dunn's test) at different time points across treatments ($P < 0.05$). See Supplementary File S3 for data and statistical analyses. **(C)** Shape coefficient (SC) of cells from control (blue) and PEG-grown (orange) calli ($n = 22$) at 6 months of mean \pm S. D. (* indicates $P < 0.05$, Welch Two Sample t-test). See Supplementary File S4 for data and statistical analyses. **(D)** Images of cells from PEG and control cultures at 6 months of culture (scale bars PEG = 100 μm , Control 200 μm)




Figure 5. Comparisons between control and 10% PEG treatments after six months in vitro culture in terms of: **(A)** medium and callus osmolarity, **(B)** proline content, **(C)** soluble sugars, and **(D)** water content. Note different letters indicate significant differences ($P \leq 0.05$) between treatments ($n=3 \pm SD$); ns = non-significant. (Kruskal Wallis followed by Dunn's test (A); Wilcoxon signed rank test (B) and Welch Two Sample t-test (C and D)). See Supplementary File S5 for data and statistical analyses.




Figure 6. Gene expression after 6 months culture on 10% PEG medium and control (MANA medium) with leaf as reference: (A) *MtSOS1*, (B) *MtWEE1*, (C) *MtSERK1*, (D) *MtP5CS* and (E) *MtCCS52*. Different letters indicate significant differences amongst treatments/tissues ($P \leq 0.05$; 1-way ANOVA; $n = 3 \pm \text{SD}$).

Parameters assessed	Control	Ionic (NaCl) stress	Osmotic (PEG) stress
Growth (fresh weight)			
Cell Viability (%)			
Embryogenic competence			
Chlorophyll (green tissue)			
Endoreduplication			
Mitotic index			
Medium osmolarity			
Callus osmolarity			
Water content			
Proline accumulation			
Soluble sugars			
Relative expression of <i>MtSERK1</i>			
Relative expression of <i>MtWEE1</i>			
Relative expression of <i>MtCCS52</i>			
Relative expression of <i>MtP5CS</i>			
Relative expression of <i>MtSOS1</i>			

Figure 7. Summarized effects on various parameters of imposing abiotic stress on embryogenic callus of *Medicago truncatula* following *in vitro* selection. Control callus tissues are compared with NaCl-tolerant callus (Elmaghrabi et al. 2013) and PEG-induced osmotic stress tolerant callus tissues in this study. Blue colour indicates increase/upregulation. Red colour indicates decrease/downregulation. When non significant compared to controls white is used.

Figure 1.JPEG




Figure 1. Mean (\pm S.D.) callus growth over 6 months on MANA medium with 10% w/v PEG compared with control (MANA medium). Different letter combinations indicate significant differences ($P \leq 0.05$; Kruskal Wallis followed by a Dunn's test) ($n > 8$). See Supplementary File S1 for data and statistical analyses.




Figure 2. Callus phenotype and viability after 6 months on 10% w/v PEG and control media: **(A)** control callus; **(B)** PEG-selected callus, arrows indicate somatic embryos; viability of control callus after FDA staining observed under transmission **(C)** or UV **(D)** light; viability of PEG-selected callus after FDA staining observed under transmission **(E)** or UV **(F)** light. Scale bars are A = 2.13 mm, B = 3.34 mm; and C through to F = 200 μm.

Figure 3.JPEG




Figure 3. FCM profile and % distribution of nuclei in G1 (1st peak), G2 (2nd peak), S phase (trough between 1st and 2nd peaks) and polyploid nuclei (3rd (8C) and 4th (16C) peaks) from (A) leaves of *Medicago truncatula*, (B) *M. truncatula* callus cultured on 10% w/v PEG6000 for 6 months; (C) Mean (\pm S.D.) mitotic index in leaves compared to callus tissues after 9 continuous months growth on 10% w/v PEG. Different letters indicate significant differences ($P \leq 0.05$); $n=10$. See Supplementary File S2 for data and statistical analyses.




Figure 4. Effects of PEG6000 on cell morphology. **(A)** cell and **(B)** nuclear size (μm^2) at different time points during *in vitro* selection for PEG6000 (10%) resistance. Data are the means \pm S.D. from $n = 9$ replicates at 2 and 4 months of culture and $n = 22$ measurements at 6 months of culture. Bars with different letters were significantly different (Kruskal Wallis followed by Dunn's test) at different time points across treatments ($P < 0.05$). See Supplementary File S3 for data and statistical analyses. **(C)** Shape coefficient (SC) of cells from control (blue) and PEG-grown (orange) calli ($n = 22$) at 6 months of mean \pm S. D. (* indicates $P < 0.05$, Welch Two Sample t-test). See Supplementary File S4 for data and statistical analyses. **(D)** Images of cells from PEG and control cultures at 6 months of culture (scale bars PEG = 100 μm , Control 200 μm)




Figure 5. Comparisons between control and 10% PEG treatments after six months in vitro culture in terms of: **(A)** medium and callus osmolarity, **(B)** proline content, **(C)** soluble sugars, and **(D)** water content. Note different letters indicate significant differences ($P \leq 0.05$) between treatments ($n=3 \pm SD$); ns = non-significant. (Kruskal Wallis followed by Dunn's test (A); Wilcoxon signed rank test (B) and Welch Two Sample t-test (C and D)). See Supplementary File S5 for data and statistical analyses.

Figure 6.JPEG




Figure 6. Gene expression after 6 months culture on 10% PEG medium and control (MANA medium) with leaf as reference: (A) *MtSOS1*, (B) *MtWEE1*, (C) *MtSERK*, (D) *MtP5CS* and (E) *MtCCS52*. Different letters indicate significant differences amongst treatments/tissues ($P \leq 0.05$; 1-way ANOVA; $n = 3 \pm \text{SD}$).

Parameters assessed	Control	Ionic (NaCl) stress	Osmotic (PEG) stress
Growth (fresh weight)			
Cell Viability (%)			
Embryogenic competence			
Chlorophyll (green tissue)			
Endoreduplication			
Mitotic index			
Medium osmolarity			
Callus osmolarity			
Water content			
Proline accumulation			
Soluble sugars			
Relative expression of <i>MtSERK1</i>			
Relative expression of <i>MtWEE1</i>			
Relative expression of <i>MtCCS52</i>			
Relative expression of <i>MtP5CS</i>			
Relative expression of <i>MtSOS1</i>			

Figure 7. Summarized effects on various parameters of imposing abiotic stress on embryogenic callus of *Medicago truncatula* following *in vitro* selection. Control callus tissues are compared with NaCl-tolerant callus (Elmaghrabi et al. 2013) and PEG-induced osmotic stress tolerant callus tissues in this study. Blue colour indicates increase/upregulation. Red colour indicates decrease/downregulation. When non significant compared to controls white is used.